HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TECVAYLI safely and effectively. See full prescribing information for TECVAYLI.

TECVAYLI® (teclistamab-cqyv) injection, for subcutaneous use Initial U.S. Approval: 2022

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

See full prescribing information for complete boxed warning.

Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving TECVAYLI. Initiate treatment with TECVAYLI step-up dosing schedule to reduce risk of CRS. Withhold TECVAYLI until CRS resolves or permanently discontinue based on severity. (2.1, 2.4, 5.1)

Neurologic toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) and serious, life-threatening or fatal reactions, can occur in patients receiving TECVAYLI. Monitor patients for signs or symptoms of neurologic toxicity, including ICANS, during treatment. Withhold TECVAYLI until neurologic toxicity resolves or permanently discontinue based on severity. (2.4, 5.2)

TECVAYLI is available only through a restricted program called the TECVAYLI and TALVEY Risk Evaluation and Mitigation Strategy (REMS). (5.3)

Boxed Warnings 8/2025 Dosage and Administration, Recommended Dosage (2.1) 2/2024 Dosage and Administration, Restarting TECVAYLI after Dosage Delay (2.3) 11/2024 Dosage and Administration, Preparation and Administration (2.5) 2/2024 Warnings and Precautions, Neurologic Toxicity including ICANS (5.2) 8/2025

TECVAYLI is a bispecific B-cell maturation antigen (BCMA)-directed CD3 T-cell engager indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody.

This indication is approved under accelerated approval based on response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s). (1)

-----DOSAGE AND ADMINISTRATION------DOSAGE AND ADMINISTRATION

TECVAYLI Recommended Dosing Schedule (2.1)				
Dosing Schedule	Day Dose			
All Patients				
	Day 1	Step-up dose 1	0.06 mg/kg	
Step-up Dosing	Day 4ª	Step-up dose 2	0.3 mg/kg	
Schedule	Day 7 ^b	First treatment dose	1.5 mg/kg	
Weekly Dosing Schedule	One week after first treatment dose and weekly thereafter	Subsequent treatment doses	1.5 mg/kg once weekly	

Patients who have achieved and maintained a complete response or better for a minimum of 6 months

Biweekly (every two weeks)			
two weeks)			
docina echadula			

2

The dosing frequency may be decreased to 1.5 mg/kg every two weeks

TECVAYLI® (teclistamab-cqyv) injection

- ^a Step-up dose 2 may be given between 2 to 4 days after step-up dose 1 and may be given up to 7 days after step-up dose 1 to allow for resolution of adverse reactions.
- b First treatment dose may be given between 2 to 4 days after step-up dose 2 and may be given up to 7 days after step-up dose 2 to allow for resolution of adverse reactions.
- For subcutaneous injection only. (2.1)
- Patients should be hospitalized for 48 hours after administration of all doses within the TECVAYLI step-up dosing schedule. (2.1)
- Administer pretreatment medications as recommended. (2.2)
- Refer to Tables 7, 8, and 9 to determine the dosage based on predetermined weight ranges. (2.5)
- See Full Prescribing Information for instructions on preparation and administration. (2.5)

----- DOSAGE FORMS AND STRENGTHS ------

Injection

- 30 mg/3 mL (10 mg/mL) in a single-dose vial (3)
- 153 mg/1.7 mL (90 mg/mL) in a single-dose vial (3)

------ CONTRAINDICATIONS -----

None (4

-----WARNINGS AND PRECAUTIONS -----

- Hepatotoxicity: Can cause hepatotoxicity, including fatalities. Monitor liver enzymes and bilirubin at baseline and during treatment as clinically indicated. (5.4)
- Infections: Can cause severe, life-threatening, or fatal infections. Monitor
 patients for signs and symptoms of infection and treat appropriately. Withhold
 in patients with active infection during the step-up dosing schedule. (2.4, 5.5)
- <u>Neutropenia</u>: Monitor complete blood cell counts at baseline and periodically during treatment. (5.6)
- <u>Hypersensitivity and Other Administration Reactions</u>: Systemic administrationrelated reactions and local injection site reactions can occur. Withhold or consider permanent discontinuation based on severity. (2.4, 5.7)
- Embryo-Fetal Toxicity: May cause fetal harm. Advise females of reproductive potential of the potential risk to the fetus and to use effective contraception. (5.8, 8.1, 8.3)

------ ADVERSE REACTIONS ------

The most common adverse reactions (\ge 20%) are pyrexia, cytokine release syndrome, musculoskeletal pain, injection site reaction, fatigue, upper respiratory tract infection, nausea, headache, pneumonia, and diarrhea.

The most common Grade 3 to 4 laboratory abnormalities (\geq 20%) are decreased lymphocytes, decreased neutrophils, decreased white blood cells, decreased hemoglobin, and decreased platelets. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Janssen Biotech, Inc. at 1-800-526-7736 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- USE IN SPECIFIC POPULATIONS -----

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 8/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: CYTOKINE RELEASE SYNDROME AND NEUROLOGIC TOXICITY INCLUDING IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

- I INDICATIONS AND USAGE
 - DOSAGE AND ADMINISTRATION
 - 2.1 Recommended Dosage
 - 2.2 Recommended Pretreatment Medications
 - 2.3 Restarting TECVAYLI after Dosage Delay
 - 2.4 Dosage Modifications for Adverse Reactions
 - 2.5 Preparation and Administration
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Cytokine Release Syndrome
 - 5.2 Neurologic Toxicity including ICANS
 - 5.3 TECVAYLI and TALVEY REMS5.4 Hepatotoxicity
 - 5.5 Infections
 - 5.6 Neutropenia
 - 5.7 Hypersensitivity and Other Administration Reactions
 - 5.8 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 7 DRUG INTERACTIONS
- B USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.3 Females and Males of Reproductive Potential
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
 - 12.6 Immunogenicity
- 13 NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES
- 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

^{*}Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving TECVAYLI. Initiate treatment with TECVAYLI step-up dosing schedule to reduce risk of CRS. Withhold TECVAYLI until CRS resolves or permanently discontinue based on severity [see Dosage and Administration (2.1, 2.4) and Warnings and Precautions (5.1)].

Neurologic toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) and serious, life-threatening or fatal reactions, can occur with TECVAYLI. Monitor patients for signs or symptoms of neurologic toxicity, including ICANS. during treatment. Withhold TECVAYLI until neurologic toxicity resolves or permanently discontinue based on severity [see Dosage and Administration (2.4) and Warnings and Precautions (5.2)].

Because of the risk of CRS and neurologic toxicity, including ICANS, TECVAYLI is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the TECVAYLI and TALVEY REMS [see Warnings and Precautions (5.3)].

INDICATIONS AND USAGE

TECVAYLI is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antihody.

This indication is approved under accelerated approval based on response rate [see Clinical Studies (14)]. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

For subcutaneous injection only.

The recommended dosing schedule for TECVAYLI is provided in Table 1. The recommended dosage of TECVAYLI is step-up doses of 0.06 mg/kg and 0.3 mg/kg followed by 1.5 mg/kg once weekly until disease progression or unacceptable toxicity. In patients who have achieved and maintained a complete response or better for a minimum of 6 months, the dosing frequency may be decreased to 1.5 mg/kg every two weeks until disease progression or unacceptable toxicity.

Administer pretreatment medications prior to each dose of the TECVAYLI stepup dosing schedule, which includes step-up dose 1, step-up dose 2, and the first treatment dose as described in Table 1 [see Dosage and Administration (2.2)].

Administer TECVAYLI subcutaneously according to the step-up dosing schedule in Table 1 to reduce the incidence and severity of cytokine release syndrome (CRS). Due to the risk of CRS and neurologic toxicity, including ICANS, patients should be hospitalized for 48 hours after administration of all doses within the TECVAYLI step-up dosing schedule [see Dosage and Administration (2.4) and Warnings and Precautions (5.1, 5.2)].

Table 1: TECVAYLI Dosing Schedule

Dosing schedule Day		Dose				
All Patients	All Patients					
	Day 1	Step-up dose 1	0.06 mg/kg			
Step-up dosing	Day 4 ^b	Step-up dose 2	0.3 mg/kg			
scheduleª	Day 7°	First treatment dose	1.5 mg/kg			
Weekly dosing schedule ^a	One week after first treatment dose and weekly thereafter	Subsequent treatment doses	1.5 mg/kg once weekly			

Patients who have achieved and maintained a complete response or better for a minimum of 6 months

Biweekly (every two weeks) dosing schedule ^a	The dosing frequency may every two

be decreased to 1.5 mg/kg

- ^a See Table 2 for recommendations on restarting TECVAYLI after dose delays [see Dosage and Administration (2.3)].
- ^b Step-up dose 2 may be given between 2 to 4 days after step-up dose 1 and may be given up to 7 days after step-up dose 1 to allow for resolution of adverse reactions.
- ^c First treatment dose may be given between 2 to 4 days after step-up dose 2 and may be given up to 7 days after step-up dose 2 to allow for resolution of adverse reactions.

Refer to Tables 7, 8, and 9 to determine the dosage based on predetermined weight ranges [see Dosage and Administration (2.5)].

2.2 Recommended Pretreatment Medications

Administer the following pretreatment medications 1 to 3 hours before each dose of the TECVAYLI step-up dosing schedule, which includes step-up dose 1, step-up dose 2, and the first treatment dose (see Table 1), to reduce the risk of CRS Isee Warnings and Precautions (5.1) and Adverse Reactions (6.1)].

- Corticosteroid (oral or intravenous dexamethasone 16 mg)
- Histamine-1 (H1) receptor antagonist (oral or intravenous diphenhydramine 50 mg or equivalent)
- Antipyretics (oral or intravenous acetaminophen 650 mg to 1,000 mg or equivalent) Administration of pretreatment medications may be required prior to administration of subsequent doses of TECVAYLI in the following patients:
- Patients who repeat doses within the TECVAYLI step-up dosing schedule following a dose delay [see Dosage and Administration (2.3)]
- Patients who experienced CRS following the prior dose of TECVAYLI [see Dosage and Administration (2.4)].

Prophylaxis for Herpes Zoster Reactivation

Prior to starting treatment with TECVAYLI, consider initiation of antiviral prophylaxis to prevent herpes zoster reactivation per guidelines.

2.3 Restarting TECVAYLI after Dosage Delay

If a dose of TECVAYLI is delayed, restart therapy based on the recommendations in Table 2 and resume the treatment schedule accordingly [see Dosage and Administration (2.1)]. Administer pretreatment medications as indicated in Table 2. Due to the risk of CRS and neurologic toxicity, including ICANS, patients should be hospitalized for 48 hours after administration of all doses within the TECVAYLI step-up dosing schedule [see Dosage and Administration (2.4) and Warnings and Precautions (5.1, 5.2)].

Table 2: Recommendations for Restarting Therapy with TECVAYLI After Dose Delay

Last dose administered	Time since the last dose administered	Action
Step-up dose 1	More than 7 days	Restart TECVAYLI step-up dosing schedule at step-up dose 1 (0.06 mg/kg).ª
Cton un dogo 2	8 days to 28 days	Repeat step-up dose 2 (0.3 mg/kg) ^a and continue TECVAYLI step-up dosing schedule.
Step-up dose 2	More than 28 days ^b	Restart TECVAYLI step-up dosing schedule at step-up dose 1 (0.06 mg/kg).ª
	28 days or less	Continue TECVAYLI at last treatment dose in weekly schedule (1.5 mg/kg once weekly; see Table 1).
Any weekly treatment dose	29 days to 56 days ^b	Restart TECVAYLI step-up dosing schedule at step-up dose 2 (0.3 mg/kg).ª
	More than 56 days ^b	Restart TECVAYLI step-up dosing schedule at step-up dose 1 (0.06 mg/kg).ª
Any biweekly	63 days or less ^b	Continue TECVAYLI at last treatment dose in biweekly schedule (1.5 mg/kg every two weeks; see Table 1).
(every two weeks) treatment dose	64 days to 112 days ^b	Restart TECVAYLI step-up dosing schedule at step up dose 2 (0.3 mg/kg). ^a
a calliont door	More than 112 days ^b	Restart TECVAYLI step-up dosing schedule at step-up dose 1 (0.06 mg/kg). ^a

^a Administer pretreatment medications prior to TECVAYLI dose and monitor patients accordingly [see Dosage and Administration (2.2, 2.5)].

^b Consider benefit-risk of restarting TECVAYLI in patients who require a dose delay of more than 28 days due to an adverse reaction.

2.4 Dosage Modifications for Adverse Reactions

Dosage reductions of TECVAYLI are not recommended.

Dosage delays may be required to manage toxicities related to TECVAYLI [see Warnings and Precautions (5)1.

See Tables 3. 4. and 5 for recommended actions for adverse reactions of CRS. neurologic toxicity, and ICANS. See Table 6 for recommended actions for other adverse reactions following administration of TECVAYLI.

Management of CRS, Neurologic Toxicity and ICANS

Cytokine Release Syndrome (CRS)

Management recommendations for CRS are summarized in Table 3.

Identify CRS based on clinical presentation [see Warnings and Precautions (5.1)]. Evaluate and treat other causes of fever, hypoxia, and hypotension.

If CRS is suspected, withhold TECVAYLI until CRS resolves. Manage according to the recommendations in Table 3 and consider further management per current practice guidelines. Administer supportive therapy for CRS, which may include intensive care for severe or life-threatening CRS. Consider laboratory testing to monitor for disseminated intravascular coagulation (DIC), hematology parameters, as well as pulmonary, cardiac, renal, and hepatic function.

Table 3: Recommendations for Management of Cytokine Release Syndrome

Gradea	Presenting Symptoms	Actions
Grade 1	Temperature	Withhold TECVAYLI until CRS resolves.
	≥100.4 °F (38 °C) ^b	Administer pretreatment medications prior to next dose of TECVAYLI.
Grade 2	Temperature ≥100.4 °F (38 °C) ^b with: Hypotension responsive to fluids and not requiring vasopressors, and/or, Oxygen requirement of low-flow nasal	Withhold TECVAYLI until CRS resolves. Administer pretreatment medications prior to next dose of TECVAYLI. Patients should be hospitalized for 48 hours following the next dose of TECVAYLI [see Dosage and Administration (2.1)].
Grade 3	cannula ^d or blow-by. Temperature ≥100.4 °F (38 °C) ^b with: Hypotension requiring one vasopressor with or without vasopressin, and/or, Oxygen requirement of high-flow nasal cannula ^d , facemask, non-rebreather mask, or Venturi mask.	First Occurrence of Grade 3 CRS with Duration Less than 48 Hours: Withhold TECVAYLI until CRS resolves. Provide supportive therapy, which may include intensive care. Administer pretreatment medications prior to next dose of TECVAYLI. Patients should be hospitalized for 48 hours following the next dose of TECVAYLI [see Dosage and Administration (2.1)]. Recurrent Grade 3 CRS or Grade 3 CRS with Duration 48 Hours or Longer: Permanently discontinue TECVAYLI. Provide supportive therapy, which may include intensive care.
Grade 4	Temperature ≥100.4 °F (38 °C) ^b with: Hypotension requiring multiple vasopressors (excluding vasopressin), and/or, Oxygen requirement of positive pressure (e.g., continuous positive airway pressure (CPAP), bilevel positive airway pressure (BiPAP), intubation, and mechanical ventilation).	Permanently discontinue TECVAYLI. Provide supportive therapy, which may include intensive care.

- ^a Based on American Society for Transplantation and Cellular Therapy (ASTCT) 2019 grading for CRS.
- b Attributed to CRS. Fever may not always be present concurrently with hypotension or hypoxia as it may be masked by interventions such as antipyretics or anticytokine therapy.
- ^c See Table 2 for recommendations on restarting TECVAYLI after dose delays [see Dosage and Administration (2.3)].
- d Low-flow nasal cannula is ≤6 L/min, and high-flow nasal cannula is >6 L/min.

Neurologic Toxicity and ICANS

Management recommendations for neurologic toxicity and ICANS are summarized in Tables 4 and 5.

At the first sign of neurologic toxicity, including ICANS, withhold TECVAYLI and consider neurology evaluation. Rule out other causes of neurologic symptoms. Provide supportive therapy, which may include intensive care, for severe or life-threatening neurologic toxicities, including ICANS [see Warnings and Precautions (5.2)]. Manage ICANS according to the recommendations in Table 5 and consider further management per current practice guidelines.

Table 4: Recommendations for Management of Neurologic Toxicity (excluding ICANS)

Adverse Reaction	Severitya	Actions
Neurologic Toxicity ^a (excluding ICANS)	Grade 1	Withhold TECVAYLI until neurologic toxicity symptoms resolve or stabilize. ^b
	Grade 2 Grade 3 (First occurrence)	Withhold TECVAYLI until neurologic toxicity symptoms improve to Grade 1 or less. ^b Provide supportive therapy.
	Grade 3 (Recurrent) Grade 4	Permanently discontinue TECVAYLI. Provide supportive therapy, which may include intensive care.

- ^a Based on National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE), version 4.03.
- ^b See Table 2 for recommendations on restarting TECVAYLI after dose delays [see Dosage and Administration (2.3)].

Table 5: Recommendations for Management of Immune Effector Cell-Associated Neurotoxicity Syndrome

Gradea	Presenting Symptoms ^b	Actions
Grade 1	ICE score 7-9°, or depressed level of	Withhold TECVAYLI until ICANS resolves. ^e
	consciousness ^d : awakens spontaneously.	Monitor neurologic symptoms and consider consultation with neurologist and other specialists for further evaluation and management, including consideration for starting non- sedating, anti-seizure medicines for seizure prophylaxis.
Grade 2	ICE score 3-6°, or depressed level of	Withhold TECVAYLI until ICANS resolves.
	consciousness ^d : awakens to voice.	Administer dexamethasone ^f 10 mg intravenously every 6 hours. Continue dexamethasone use until resolution to Grade 1 or less then taper.
		Monitor neurologic symptoms and consider consultation with neurologist and other specialists for further evaluation and management, including consideration for starting non- sedating, anti-seizure medicines for seizure prophylaxis.
		Patients should be hospitalized for 48 hours following the next dose of TECVAYLI [see Dosage and Administration (2.1)].

Table 5: Recommendations for Management of Immune Effector Cell-Associated Neurotoxicity Syndrome (continued)

Gradea	Presenting Symptoms ^b	Actions
Grade 3	ICE score 0-2°.	First Occurrence of Grade 3 ICANS:
Graue 3	or depressed level of consciousness ^d : awakens	Withhold TECVAYLI until ICANS resolves.
	only to tactile stimulus, or seizures ^d , either: • any clinical seizure, focal or generalized,	 Administer dexamethasone^f 10 mg intravenously every 6 hours. Continue dexamethasone use until resolution to Grade 1 or less, then taper.
	that resolves rapidly, or non-convulsive seizures on electroencephalogram (EEG) that resolve with intervention,	Monitor neurologic symptoms and consider consultation with neurologist and other specialists for further evaluation and management, including consideration for starting non- sedating, anti-seizure medicines for seizure prophylaxis.
	or raised intracranial pressure: focal/local	 Provide supportive therapy, which may include intensive care.
	edema on neuroimaging ^d .	 Patients should be hospitalized for 48 hours following the next dose of TECVAYLI [see Dosage and Administration (2.1)].°
		Recurrent Grade 3 ICANS:
		Permanently discontinue TECVAYLI
		 Administer dexamethasone^f 10 mg intravenously and repeat dose every 6 hours. Continue dexamethasone use until resolution to Grade 1 or less, then taper.
		 Monitor neurologic symptoms and consider consultation with neurologist and other specialists for further evaluation and management, including consideration for starting non- sedating, anti-seizure medicines for seizure prophylaxis.
		Provide supportive therapy, which may include intensive care.

Table 5: Recommendations for Management of Immune Effector Cell-Associated Neurotoxicity Syndrome (continued)

Neurotoxicity Syndrome (continued)			
Gradea	Presenting Symptoms ^b		Actions
Grade 4	ICE score 0°,	•	Permanently discontinue TECVAYLI.
	or depressed level of consciousness ^d : either:	•	Administer dexamethasone ^f 10 mg intravenously and repeat dose every 6
	 patient is unarousable or requires vigorous or repetitive tactile stimuli to arouse, or 	•	hours. Continue dexamethasone use until resolution to Grade 1 or less, then taper. Alternatively, consider administration
	 stupor or coma, 		of methylprednisolone 1,000 mg
	or seizures ^d , either:		per day intravenously and continue methylprednisolone 1,000 mg per day
	 life-threatening 		intravenously for 2 or more days.
	prolonged seizure (>5 minutes), or	•	Monitor neurologic symptoms and consider consultation with neurologist
	 repetitive clinical or electrical seizures without return to baseline in between, 	and other specialists for further evaluation and management, inclu consideration for starting non-	evaluation and management, including consideration for starting non- sedating, anti-seizure medicines for
	or motor findings ^d :		seizure prophylaxis.
	 deep focal motor weakness such as hemiparesis or paraparesis, 	•	Provide supportive therapy, which may include intensive care.
	or raised intracranial pressure/cerebral edema ^d , with signs/ symptoms such as:		
	 diffuse cerebral edema on neuroimaging, or 		
	 decerebrate or decorticate posturing, or 		
	 cranial nerve VI palsy, or 		
	 papilledema, or 		
	• Cushing's triad.		

- ^a Based on American Society for Transplantation and Cellular Therapy (ASTCT) 2019 grading for ICANS.
- b Management is determined by the most severe event, not attributable to any other cause.
- c If patient is arousable and able to perform Immune Effector Cell-Associated Encephalopathy (ICE) Assessment, assess: **Orientation** (oriented to year, month, city, hospital = 4 points); **Naming** (name 3 objects, e.g., point to clock, pen, button = 3 points); **Following Commands** (e.g., "show me 2 fingers" or "close your eyes and stick out your tongue" = 1 point); **Writing** (ability to write a standard sentence = 1 point; and **Attention** (count backwards from 100 by ten = 1 point). If patient is unarousable and unable to perform ICE Assessment (Grade 4 ICANS) = 0 points.
- d Not attributable to any other cause.
- ^e See Table 2 for recommendations on restarting TECVAYLI after dose delays [see Dosage and Administration (2.3)].
- ^f All references to dexamethasone administration are dexamethasone or equivalent.

Table 6: Recommended Dosage Modifications for Other Adverse Reactions

Adverse	Severity		Actions
Infections ^a [see Warnings and Precautions	All Grades	•	Withhold TECVAYLI in patients with active infection during the step-up dosing schedule. ^b
(5.5)]	Grade 3	•	Withhold subsequent treatment doses of TECVAYLI (i.e., doses administered after TECVAYLI step-up dosing schedule) until infection improves to Grade 1 or less. ^b
	Grade 4	•	Consider permanent discontinuation of TECVAYLI.
		•	If TECVAYLI is not permanently discontinued, withhold subsequent treatment doses of TECVAYLI (i.e., doses administered after TECVAYLI step-up dosing schedule) until infection improves to Grade 1 or less. ^b
Hematologic Toxicities [see Warnings and Precautions (5.6) and Adverse Reactions (6.1)]	Absolute neutrophil count less than $0.5 \times 10^9/L$	•	Withhold TECVAYLI until absolute neutrophil count is $0.5 \times 10^9/L$ or higher. ^b
	Febrile neutropenia	•	Withhold TECVAYLI until absolute neutrophil count is 1 × 10 ⁹ /L or higher and fever resolves. ^b
	Hemoglobin less than 8 g/dL	•	Withhold TECVAYLI until hemoglobin is 8 g/dL or higher. ^b
	Platelet count less than 25,000/mcL Platelet count between 25,000/mcL and 50,000/mcL with bleeding	•	Withhold TECVAYLI until platelet count is 25,000/mcL or higher and no evidence of bleeding. ^b
Other Non- Hematologic	Grade 3	•	Withhold TECVAYLI until adverse reaction improves to Grade 1 or less. ^b
Adverse Reactions ^a	Grade 4	•	Consider permanent discontinuation of TECVAYLI.
[see Warnings and Precautions (5.4) and Adverse Reactions (6.1)]		•	If TECVAYLI is not permanently discontinued, withhold subsequent treatment doses of TECVAYLI (i.e., doses administered after TECVAYLI step-up dosing schedule) until adverse reaction improves to Grade 1 or less. ^b

^a Based on National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE), version 4.03.

2.5 Preparation and Administration

TECVAYLI is intended for subcutaneous use by a healthcare provider only.

TECVAYLI should be administered by a healthcare provider with adequate medical personnel and appropriate medical equipment to manage severe reactions, including CRS and ICANS [see Warnings and Precautions (5.1, 5.2)].

TECVAYLI is a clear to slightly opalescent, colorless to light yellow solution. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use if the solution is discolored, or cloudy, or if foreign particles are present.

TECVAYLI 30 mg/3 mL (10 mg/mL) vial and TECVAYLI 153 mg/1.7 mL (90 mg/mL) vial are supplied as ready-to-use solution that do not need dilution prior to administration.

Do not combine TECVAYLI vials of different concentrations to achieve treatment dose.

Use aseptic technique to prepare and administer TECVAYLI.

Preparation of TECVAYLI

Refer to the following reference tables for the preparation of TECVAYLI.

Refer to Tables 7, 8, and 9 below to determine the dosage based on predetermined weight ranges.

Use Table 7 to determine total dose, injection volume and number of vials required based on patient's actual body weight for step-up dose 1 using TECVAYLI 30 mg/3 mL (10 mg/mL) vial.

Table 7: Step-up Dose 1 (0.06 mg/kg) Injection Volumes using TECVAYLI 30 mg/3 mL (10 mg/mL) Vial

Patient Body Weight (kg)	Total Dose (mg)	Volume of Injection (mL)	Number of Vials (1 vial=3 mL)
35 to 39	2.2	0.22	1
40 to 44	2.5	0.25	1
45 to 49	2.8	0.28	1
50 to 59	3.3	0.33	1
60 to 69	3.9	0.39	1
70 to 79	4.5	0.45	1
80 to 89	5.1	0.51	1
90 to 99	5.7	0.57	1
100 to 109	6.3	0.63	1
110 to 119	6.9	0.69	1
120 to 129	7.5	0.75	1
130 to 139	8.1	0.81	1
140 to 149	8.7	0.87	1
150 to 160	9.3	0.93	1

Use Table 8 to determine total dose, injection volume and number of vials required based on patient's actual body weight for step-up dose 2 using TECVAYLI 30 mg/3 mL (10 mg/mL) vial.

Table 8: Step-up Dose 2 (0.3 mg/kg) Injection Volumes using TECVAYLI 30 mg/3 mL (10 mg/mL) Vial

(10 1119/1112) 1141			
Patient Body Weight (kg)	Total Dose (mg)	Volume of Injection (mL)	Number of Vials (1 vial=3 mL)
35 to 39	11	1.1	1
40 to 44	13	1.3	1
45 to 49	14	1.4	1
50 to 59	16	1.6	1
60 to 69	19	1.9	1
70 to 79	22	2.2	1
80 to 89	25	2.5	1
90 to 99	28	2.8	1
100 to 109	31	3.1	2
110 to 119	34	3.4	2
120 to 129	37	3.7	2
130 to 139	40	4	2
140 to 149	43	4.3	2
150 to 160	47	4.7	2

Use Table 9 to determine total dose, injection volume and number of vials required based on patient's actual body weight for the treatment dose using TECVAYLI 153 mg/1.7 mL (90 mg/mL) vial.

Table 9: Treatment Dose (1.5 mg/kg) Injection Volumes using TECVAYLI 153 mg/1.7 mL (90 mg/mL) Vial

(30 Hig/IIIL/ Viai			
Patient Body Weight (kg)	Total Dose (mg)	Volume of Injection (mL)	Number of Vials (1 vial=1.7 mL)
35 to 39	56	0.62	1
40 to 44	63	0.7	1
45 to 49	70	0.78	1
50 to 59	82	0.91	1
60 to 69	99	1.1	1
70 to 79	108	1.2	1
80 to 89	126	1.4	1
90 to 99	144	1.6	1
100 to 109	153	1.7	1
110 to 119	171	1.9	2
120 to 129	189	2.1	2
130 to 139	198	2.2	2
140 to 149	216	2.4	2
150 to 160	234	2.6	2

^b See Table 2 for recommendations on restarting TECVAYLI after dose delays [see Dosage and Administration (2.3)].

Remove the appropriate strength TECVAYLI vial from refrigerated storage [2 °C to 8 °C (36 °F to 46 °F)].

Once removed from refrigerated storage, equilibrate TECVAYLI to ambient temperature [15 °C to 30 °C (59 °F to 86 °F)] for at least 15 minutes. Do not warm TECVAYLI in any other way.

Once equilibrated, gently swirl the vial for approximately 10 seconds to mix. Do not shake.

Withdraw the required injection volume of TECVAYLI from the vial(s) into an appropriately sized syringe using a transfer needle.

Each injection volume should not exceed 2 mL. Divide doses requiring greater than 2 mL equally into multiple syringes.

TECVAYLI is compatible with stainless steel injection needles and polypropylene or polycarbonate syringe material.

Replace the transfer needle with an appropriately sized needle for injection.

Administration of TECVAYLI

Inject the required volume of TECVAYLI into the subcutaneous tissue of the abdomen (preferred injection site). Alternatively, TECVAYLI may be injected into the subcutaneous tissue at other sites (e.g., thigh). If multiple injections are required, TECVAYLI injections should be at least 2 cm apart.

Do not inject into tattoos or scars or areas where the skin is red, bruised, tender, hard or not intact.

Any unused product or waste material should be disposed in accordance with local requirements.

<u>Storage</u>

If the prepared dosing syringe(s) of TECVAYLI is not used immediately, store syringe(s) at 2 °C to 8 °C (36 °F to 46 °F) or at ambient temperature 15 °C to 30 °C (59 °F to 86 °F) for a maximum of 20 hours. Discard syringe(s) after 20 hours, if not used.

Monitoring

Due to the risk of CRS and neurologic toxicity, including ICANS, patients should be hospitalized for 48 hours after administration of all doses within the TECVAYLI step-up dosing schedule [see Dosage and Administration (2.1) and Warnings and Precautions (5.1, 5.2)].

3 DOSAGE FORMS AND STRENGTHS

Injection

- 30 mg/3 mL (10 mg/mL) clear to slightly opalescent, colorless to light yellow solution in a single-dose vial.
- 153 mg/1.7 mL (90 mg/mL) clear to slightly opalescent, colorless to light yellow solution in a single-dose vial.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Cytokine Release Syndrome

TECVAYLI can cause cytokine release syndrome (CRS), including life-threatening or fatal reactions [see Adverse Reactions (6.1)].

In the clinical trial, CRS occurred in 72% of patients who received TECVAYLI at the recommended dose, with Grade 1 CRS occurring in 50% of patients, Grade 2 in 21%, and Grade 3 in 0.6%. Recurrent CRS occurred in 33% of patients. Most patients experienced CRS following step-up dose 1 (42%), step-up dose 2 (35%), or the initial treatment dose (24%). Less than 3% of patients developed first occurrence of CRS following subsequent doses of TECVAYLI. The median time to onset of CRS was 2 (range: 1 to 6) days after the most recent dose with a median duration of 2 (range: 1 to 9) days.

Clinical signs and symptoms of CRS included, but were not limited to, fever, hypoxia, chills, hypotension, sinus tachycardia, headache, and elevated liver enzymes (aspartate aminotransferase and alanine aminotransferase elevation).

Initiate therapy according to TECVAYLI step-up dosing schedule to reduce risk of CRS [see Dosage and Administration (2.1, 2.4]]. Administer pretreatment medications to reduce risk of CRS and monitor patients following administration of TECVAYLI accordingly [see Dosage and Administration (2.2, 2.4]].

At the first sign of CRS, immediately evaluate patient for hospitalization. Administer supportive care based on severity and consider further management per current practice guidelines. Withhold or permanently discontinue TECVAYLI based on severity [see Dosage and Administration (2.4)].

TECVAYLI is available only through a restricted program under a REMS [see Warnings and Precautions (5.3)].

5.2 Neurologic Toxicity including ICANS

TECVAYLI can cause serious, life-threatening or fatal neurologic toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) [see Adverse Reactions (6.1)].

In the clinical trial, neurologic toxicity occurred in 57% of patients who received TECVAYLI at the recommended dose, with Grade 3 or 4 neurologic toxicity occurring

TECVAYLI® (teclistamab-cgyv) injection

in 2.4% of patients. The most frequent neurologic toxicities were headache (25%), motor dysfunction (16%), sensory neuropathy (15%), and encephalopathy (13%).

With longer follow-up, Grade 4 seizure and fatal Guillain-Barré syndrome (one patient each) occurred in patients who received TECVAYLI.

In the clinical trial, ICANS was reported in 6% of patients who received TECVAYLI at the recommended dose [see Adverse Reactions (6.1)]. Recurrent ICANS occurred in 1.8% of patients. Most patients experienced ICANS following step-up dose 1 (1.2%), step-up dose 2 (0.6%), or the initial treatment dose (1.8%). Less than 3% of patients developed first occurrence of ICANS following subsequent doses of TECVAYLI. The median time to onset of ICANS was 4 (range: 2 to 8) days after the most recent dose with a median duration of 3 (range: 1 to 20) days. The most frequent clinical mainfestations of ICANS reported were confusional state and dysgraphia. The onset of ICANS can be concurrent with CRS, following resolution of CRS, or in the absence of CRS.

Monitor patients for signs and symptoms of neurologic toxicity during treatment. At the first sign of neurologic toxicity, including ICANS, immediately evaluate patient and provide supportive therapy based on severity. Withhold or permanently discontinue TECVAYLI based on severity per recommendations and consider further management per current practice guidelines [see Dosage and Administration (2.4)].

Due to the potential for neurologic toxicity, patients receiving TECVAYLI are at risk of depressed level of consciousness [see Adverse Reactions (6.1)]. Advise patients to refrain from driving or operating heavy or potentially dangerous machinery during and for 48 hours after completion of TECVAYLI step-up dosing schedule and in the event of new onset of any neurologic toxicity symptoms until neurologic toxicity resolves [see Dosage and Administration (2.1)].

TECVAYLI is available only through a restricted program under a REMS [see Warnings and Precautions (5.3)].

5.3 TECVAYLI and TALVEY REMS

TECVAYLI is available only through a restricted program under a REMS called the TECVAYLI and TALVEY REMS because of the risks of CRS and neurologic toxicity, including ICANS [see Warnings and Precautions (5.1, 5.2)].

Notable requirements of the TECVAYLI and TALVEY REMS include the following:

- Prescribers must be certified with the program by enrolling and completing training.
- Prescribers must counsel patients receiving TECVAYLI about the risk of CRS and neurologic toxicity, including ICANS, and provide patients with Patient Wallet Card
- Pharmacies and healthcare settings that dispense TECVAYLI must be certified with the TECVAYLI and TALVEY REMS program and must verify prescribers are certified through the TECVAYLI and TALVEY REMS program.
- Wholesalers and distributers must only distribute TECVAYLI to certified pharmacies or healthcare settings.

Further information about the TECVAYLI and TALVEY REMS program is available at www.TEC-TALREMS.com or by telephone at 1-855-810-8064.

5.4 Hepatotoxicity

TECVAYLI can cause hepatotoxicity, including fatalities. In patients who received TECVAYLI at the recommended dose in the clinical trial, there was one fatal case of hepatic failure. Elevated aspartate aminotransferase (AST) occurred in 34% of patients, with Grade 3 or 4 elevations in 1.2%. Elevated alanine aminotransferase (ALT) occurred in 28% of patients, with Grade 3 or 4 elevations in 1.8%. Elevated total bilirubin occurred in 6% of patients with Grade 3 or 4 elevations in 0.6%. Liver enzyme elevation can occur with or without concurrent CRS.

Monitor liver enzymes and bilirubin at baseline and during treatment as clinically indicated. Withhold TECVAYLI or consider permanent discontinuation of TECVAYLI based on severity [see Dosage and Administration (2.4)].

5.5 Infections

TECVAYLI can cause severe, life-threatening, or fatal infections. In patients who received TECVAYLI at the recommended dose in the clinical trial, serious infections, including opportunistic infections, occurred in 30% of patients, with Grade 3 or 4 infections in 35%, and fatal infections in 4.2% [see Adverse Reactions (6.1)].

Monitor patients for signs and symptoms of infection prior to and during treatment with TECVAYLI and treat appropriately. Administer prophylactic antimicrobials according to guidelines [see Dosage and Administration (2.2)].

Withhold TECVAYLI or consider permanent discontinuation of TECVAYLI based on severity [see Dosage and Administration (2.4)].

Monitor immunoglobulin levels during treatment with TECVAYLI and treat according to guidelines, including infection precautions and antibiotic or antiviral prophylaxis [see Dosage and Administration (2.2)].

5.6 Neutropenia

TECVAYLI can cause neutropenia and febrile neutropenia. In patients who received TECVAYLI at the recommended dose in the clinical trial, decreased neutrophils occurred in 84% of patients, with Grade 3 or 4 decreased neutrophils in 56%. Febrile neutropenia occurred in 3% of patients [see Adverse Reactions (6.1)].

Monitor complete blood cell counts at baseline and periodically during treatment and provide supportive care per local institutional guidelines.

Monitor patients with neutropenia for signs of infection.

Withhold TECVAYLI based on severity [see Dosage and Administration (2.4)].

5.7 Hypersensitivity and Other Administration Reactions

TECVAYLI can cause both systemic administration-related reactions and local injection-site reactions.

Systemic Reactions

In patients who received TECVAYLI at the recommended dose in the clinical trial, 1.2% of patients experienced systemic-administration reactions, which included Grade 1 recurrent pyrexia and Grade 1 swollen tongue.

Local Reactions

In patients who received TECVAYLI at the recommended dose in the clinical trial, injection-site reactions occurred in 35% of patients with Grade 1 injection-site reactions in 30% and Grade 2 in 4.8%.

Withhold TECVAYLI or consider permanent discontinuation of TECVAYLI based on severity [see Dosage and Administration (2.4)].

5.8 Embryo-Fetal Toxicity

Based on its mechanism of action, TECVAYLI may cause fetal harm when administered to a pregnant woman. Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with TECVAYLI and for 5 months after the last dose [see Use in Specific Populations (8.1, 8.3)].

6 ADVERSE REACTIONS

The following adverse reactions are also described elsewhere in the labeling:

- Cvtokine Release Syndrome [see Warnings and Precautions (5.1)]
- Neurologic Toxicity including ICANS [see Warnings and Precautions (5.2)]
- Hepatotoxicity [see Warnings and Precautions (5.4)]
- Infections [see Warnings and Precautions (5.5)]
- Neutropenia [see Warnings and Precautions (5.6)]
- Hypersensitivity and Other Administration Reactions [see Warnings and Precautions (5.7)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Relapsed/Refractory Multiple Myeloma

MajesTEC-1

The safety of TECVAYLI was evaluated in MajesTEC-1 [see Clinical Studies (14)] which included adult patients with relapsed or refractory multiple myeloma. Patients received step-up doses of 0.06 mg/kg and 0.3 mg/kg of TECVAYLI followed by TECVAYLI 1.5 mg/kg, subcutaneously once weekly (N=165). Among patients who received TECVAYLI, 47% were exposed for 6 months or longer and 7% were exposed for one year or longer.

The median age of patients who received TECVAYLI was 64 years (range: 33 to 84 years); 58% were male; 81% were White, 13% were Black or African American, and 2% were Asian.

Serious adverse reactions occurred in 54% of patients who received TECVAYLI. Serious adverse reactions in >2% of patients included pneumonia (15%), cytokine release syndrome (8%), sepsis (6%), general physical health deterioration (6%), COVID-19 (6%), acute kidney injury (4.8%), pyrexia (4.8%), musculoskeletal pain (2.4%), and encephalopathy (2.4%).

Fatal adverse reactions occurred in 5% of patients who received TECVAYLI, including COVID-19 (1.8%), pneumonia (1.8%), septic shock (0.6%), acute renal failure (0.6%), and hemoperitoneum (0.6%).

Permanent discontinuation of TECVAYLI due to adverse reactions occurred in 1.2% of patients. Adverse reactions resulting in permanent discontinuation of TECVAYLI included pneumonia (adenoviral and pneumocystis jirovecii pneumonia in the same patient) and hypercalcemia.

Dosage interruptions of TECVAYLI due to an adverse reaction occurred in 73% of patients. Adverse reactions which required dosage interruption in >5% of patients included neutropenia, pneumonia, pyrexia, cytokine release syndrome, upper respiratory tract infection, and COVID-19.

The most common adverse reactions (\geq 20%) were pyrexia, CRS, musculoskeletal pain, injection site reaction, fatigue, upper respiratory tract infection, nausea, headache, pneumonia, and diarrhea. The most common Grade 3 to 4 laboratory abnormalities (\geq 20%) were decreased lymphocytes, decreased neutrophils, decreased white blood cells, decreased hemoglobin, and decreased platelets.

Table 10 summarizes the adverse reactions in MajesTEC-1.

Table 10: Adverse Reactions (≥10%) in Patients with Multiple Myeloma Who Received TECVAYLI in MajesTEC-1

	(N=	TECVAYLI (N=165)	
Adverse Reactions	Any Grade (%)	Grade 3 or 4 (%)	
General disorders and administration site conditio	ns		
Pyrexia	76	3#	
Injection site reaction ¹	37	0.6#	
Fatigue ²	33	2.4#	
Chills	16	0	
Pain ³	15	1.8#	
Edema ⁴	13	0	
Immune system disorders			
Cytokine release syndrome	72	0.6#	
Hypogammaglobulinemia ⁵	11	1.2#	
Musculoskeletal and connective tissue disorde	ers		
Musculoskeletal pain ⁶	44	4.2#	
Bone pain	16	3#	
Infections			
Upper respiratory tract infection ⁷	26	2.4#	
Pneumonia ^{8*}	24	15	
Urinary tract infection ⁹	11	5#	
Gastrointestinal disorders			
Nausea	25	0.6#	
Diarrhea	21	2.4#	
Constipation	18	0	
Vomiting	12	0.6#	
Nervous system disorders			
Headache	25	0.6#	
Motor dysfunction ¹⁰	16	0	
Sensory neuropathy ¹¹	15	1.2#	
Encephalopathy ¹²	13	0	
Vascular disorders			
Hypotension	18	1.2#	
Hemorrhage ^{13*}	12	1.8	
Hypertension ¹⁴	12	4.8#	
Respiratory, thoracic, and mediastinal disorders			
Нурохіа	18	1.8	
Cough ¹⁵	15	0	
Cardiac disorders			
Cardiac arrhythmia ¹⁶	16	1.8	
Metabolism and nutrition disorders			
Decreased appetite	11	0.6#	
Renal and urinary disorders			
Acute kidney injury ¹⁷	11	3.6	

Adverse reactions were graded based on CTCAE Version 4.03, with the exception of CRS, which was graded per ASTCT 2019 criteria.

- Injection site reaction includes application site erythema, injection site bruising, injection site cellulitis, injection site discomfort, injection site erythema, injection site hematoma, injection site induration, injection site inflammation, injection site edema, injection site pruritus, injection site rash, injection site reaction and injection site swelling.
- ² Fatigue includes asthenia and fatigue
- ³ Pain includes ear pain, flank pain, groin pain, oropharyngeal pain, pain, pain in jaw, toothache and tumor pain.
- Edema includes face edema, fluid overload, fluid retention, edema peripheral and peripheral swelling.
- 5 Hypogammaglobulinemia includes hypogammaglobulinemia and hypoglobulinemia.
- Musculoskeletal pain includes arthralgia, back pain, muscle discomfort, musculoskeletal chest pain, musculoskeletal pain, myalgia, neck pain, noncardiac chest pain and pain in extremity.
- Upper respiratory tract infection includes bronchitis, influenza like illness, nasopharyngitis, pharyngitis, respiratory tract infection, respiratory tract infection bacterial, rhinitis, rhinovirus infection, sinusitis, tracheitis, upper respiratory tract infection and viral upper respiratory tract infection.
- Pneumonia includes COVID-19 pneumonia, enterobacter pneumonia, lower respiratory tract infection, metapneumovirus pneumonia, pneumocystis jirovecii pneumonia, pneumonia, pneumonia adenoviral, pneumonia klebsiella, pneumonia moraxella, pneumonia pneumococcal, pneumonia pseudomonal, pneumonia respiratory syncytial viral, pneumonia staphylococcal and pneumonia viral.

- ⁹ Urinary tract infection includes cystitis, cystitis escherichia, cystitis klebsiella, escherichia urinary tract infection, urinary tract infection and urinary tract infection bacterial.
- Motor dysfunction includes cogwheel rigidity, dysgraphia, dysphonia, gait disturbance, hypokinesia, muscle rigidity, muscle spasms, muscular weakness, peroneal nerve palsy, psychomotor hyperactivity, tremor and VIth nerve paralysis.
- Sensory neuropathy includes dysesthesia, hypoesthesia, hypoesthesia oral, neuralgia, paresthesia, paresthesia oral, peripheral sensory neuropathy, sciatica and vestibular neuronitis.
- 12 Encephalopathy includes agitation, apathy, aphasia, confusional state, delirium, depressed level of consciousness, disorientation, dyscalculia, hallucination, lethargy, memory impairment, mental status changes and somnolence.
- Hemorrhage includes conjunctival hemorrhage, epistaxis, hematoma, hematuria, hemoperitoneum, hemorrhoidal hemorrhage, lower gastrointestinal hemorrhage, melena, mouth hemorrhage and subdural hematoma.
- ¹⁴ Hypertension includes essential hypertension and hypertension.
- 15 Cough includes allergic cough, cough, productive cough and upper-airway cough syndrome.
- 16 Cardiac arrhythmia includes atrial flutter, cardiac arrest, sinus bradycardia, sinus tachycardia, supraventricular tachycardia, tachycardia and ventricular tachycardia.
- ¹⁷ Acute kidney injury includes acute kidney injury and renal impairment.
- # Only grade 3 adverse reactions occurred.
- Includes the following fatal adverse reactions: hemorrhage (n=1), pneumonia (n=3).

Clinically relevant adverse reactions in <10% of patients who received TECVAYLI included febrile neutropenia, sepsis, ICANS, seizure, Guillain-Barré syndrome, hepatic failure, and new onset or reactivated viral infections (including adenovirus, hepatitis B virus (HBV), cytomegalovirus (CMV), varicella zoster virus (VZV), herpes simplex virus (HSV), and progressive multifocal leukoencephalopathy (PML).

Table 11 summarizes laboratory abnormalities in MajesTEC-1.

Table 11: Select Laboratory Abnormalities (≥30%) That Worsened from Baseline in Patients with Multiple Myeloma Who Received TECVAYLI in MaiesTEC-1

IVIAJES I EU- I			
		TECVAYLI (N=165 ¹)	
	All Grades	Grade 3 or 4	
Laboratory Abnormality	(%)	(%)	
Hematology			
Lymphocyte count decreased	92	84	
White blood cell decreased	86	41	
Neutrophil count decreased	84	56	
Platelet count decreased	71	22	
Hemoglobin decreased	67	33	
Chemistry			
Albumin decreased	68	6	
Alkaline phosphatase increased	42	2.4	
Phosphorus decreased	38	13	
Gamma-glutamyl transferase increased	37	8	
Sodium decreased	35	10	
Aspartate aminotransferase increased	34	1.2	
Calcium (corrected) decreased	31	1.2	
Creatinine increased	30	3	

¹ The denominator used to calculate the rate varied from 164 to 165 based on the number of patients with a baseline value and at least one post-treatment value. Laboratory toxicity grades are derived based on the NCI CTCAE (National Cancer Institute Common Terminology Criteria for Adverse Events) Version 4.03.

7 DRUG INTERACTIONS

TECVAYLI causes release of cytokines [see Clinical Pharmacology (12.2)] that may suppress activity of cytochrome P450 (CYP) enzymes, resulting in increased exposure of CYP substrates. The highest risk of drug-drug interaction is expected to occur from initiation of TECVAYLI step-up dosing schedule up to 7 days after the first treatment dose and during and after CRS [see Warnings and Precautions (5.1)]. Monitor for toxicity or concentrations of drugs that are CYP substrates where minimal concentration changes may lead to serious adverse reactions. Adjust the dose of the concomitant CYP substrate drug as needed.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on the mechanism of action, TECVAYLI may cause fetal harm when administered to a pregnant woman [see Clinical Pharmacology (12.1)]. There are no available data on the use of TECVAYLI in pregnant women to evaluate

TECVAYLI® (teclistamab-cgyv) injection

for a drug associated risk. No animal reproductive or developmental toxicity studies have been conducted with TECVAYLI. Teclistamab-cqyv causes T-cell activation and cytokine release; immune activation may compromise pregnancy maintenance. Human immunoglobulin G (IgG) is known to cross the placenta; therefore, teclistamab-cqyv has the potential to be transmitted from the mother to the developing fetus. Advise women of the potential risk to the fetus.

TECVAYLI is associated with hypogammaglobulinemia, therefore, assessment of immunoglobulin levels in newborns of mothers treated with TECVAYLI should be considered.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

8.2 Lactation

Risk Summary

There are no data on the presence of teclistamab-cqyv in human milk, the effect on the breastfed child, or the effects on milk production. Maternal IgG is known to be present in human milk. The effects of local gastrointestinal exposure and limited systemic exposure in the breastfed child to TECVAYLI are unknown. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with TECVAYLI and for 5 months after the last dose.

8.3 Females and Males of Reproductive Potential

TECVAYLI may cause fetal harm when administered to a pregnant woman [see Use in Specific Populations (8.1)].

Pregnancy Testing

Verify pregnancy status of females of reproductive potential prior to initiating TECVAYLI.

Contraception

Females

Advise females of reproductive potential to use effective contraception during treatment and for 5 months after the last dose of TECVAYLI.

8.4 Pediatric Use

The safety and efficacy of TECVAYLI have not been established in pediatric patients.

8.5 Geriatric Use

Of the 165 patients with relapsed or refractory multiple myeloma treated with TECVAYLI in MajesTEC-1 at the recommended dosage, 48% were 65 years of age or older, and 15% were 75 years of age or older. No overall differences in safety or effectiveness were observed between patients 65 to 74 years of age compared to younger patients. There is an insufficient number of patients 75 years of age or older to assess whether there are differences in safety or effectiveness.

11 DESCRIPTION

Teclistamab-cqyv, a bispecific B-cell maturation antigen (BCMA)-directed CD3 T-cell engager, is a humanized immunoglobulin G4-proline, alanine, alanine (IgG4-PAA) antibody. Teclistamab-cqyv is produced in Chinese Hamster Ovary (CHO) cells using recombinant DNA technology. Teclistamab-cqyv consists of an anti-BCMA heavy chain and light chain and an anti-CD3 heavy chain and light chain with two interchain disulfide bonds connecting the two arms. The molecular weight of teclistamab-cqyv is approximately 146 kDa.

TECVAYLI® (teclistamab-cqyv) injection is a sterile, preservative-free, clear to slightly opalescent, colorless to light yellow solution supplied in a single-dose vial for subcutaneous administration.

Each TECVAYLI 3 mL single-dose vial contains 30 mg of teclistamab-cqyv, edetate disodium (0.054 mg), glacial acetic acid (0.72 mg), polysorbate 20 (1.2 mg), sodium acetate (2.7 mg), sucrose (240 mg), and Water for Injection, USP.

Each TECVAYLI 1.7 mL single-dose vial contains 153 mg of teclistamab-cqyv, edetate disodium (0.031 mg), glacial acetic acid (0.41 mg), polysorbate 20 (0.68 mg), sodium acetate (1.5 mg), sucrose (140 mg), and Water for Injection, USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Teclistamab-cqyv is a bispecific T-cell engaging antibody that binds to the CD3 receptor expressed on the surface of T-cells and B-cell maturation antigen (BCMA) expressed on the surface of multiple myeloma cells and some healthy B-lineage cells.

In vitro, teclistamab-cqyv activated T-cells, caused the release of various proinflammatory cytokines, and resulted in the lysis of multiple myeloma cells.

12.2 Pharmacodynamics

Serum concentrations of cytokines (IL-6, IL-10, TNF- α , and IFN- γ) and IL-2R were measured before and after administration of step-up dose 1, step-up dose 2, and the first three treatment doses of TECVAYLI. Increased concentrations of IL-6, IL-10, and IL-2R were observed during this period.

12.3 Pharmacokinetics

The C_{max} and AUC_{tau} of teclistamab-cqyv after the first subcutaneous treatment dose increase proportionally over a dosage range of 0.08 mg/kg to 3 mg/kg (0.05 to 2 times the approved recommended treatment dosage). Ninety percent of steady state exposure was achieved after 12 weekly treatment doses. The mean accumulation ratio between the first and 13th weekly treatment dose of teclistamab-cqyv 1.5 mg/kg was 4.2-fold for C_{max} , 4.1-fold for C_{trough} , and 5.3-fold for AUC_{tau}.

The C_{max}, C_{trough}, and AUC_{tau} of teclistamab-cqyv are presented in Table 12.

Table 12: Pharmacokinetic Parameters of Teclistamab-cqyv for the 13th Weekly
Treatment Dose of 1.5 mg/kga

ireaulient bose of 1.5 mg/kg	
Pharmacokinetic Parameter	Teclistamab-cqyv Geometric Mean (CV%)
C _{max} (mcg/mL)	23.8 (55%)
C _{trough} (mcg/mL)	21.1 (63%)
AUCtau (mcq·h/mL)	3,838 (57%)

SD = standard deviation; C_{max} = Maximum serum teclistamab-cqyv concentration; C_{trough} = Serum teclistamab-cqyv concentration prior to next dose; CV = geometric coefficient of variation; AUC_{tau} = Area under the concentration-time curve over the weekly dosing interval.

^a Following administration of teclistamab-cqyv in patients with relapsed or refractory multiple myeloma (MaiesTEC-1).

Absorption

The mean bioavailability of teclistamab-cqyv was 72% when administered subcutaneously. The median (range) T_{max} of teclistamab-cqyv after the first and 13th weekly treatment doses were 139 (19 to 168) hours and 72 (24 to 168) hours, respectively.

Distribution

The mean (coefficient of variation [CV]%) volume of distribution of teclistamab-cqyv was 5.63 L (29%).

Elimination

Teclistamab-cqyv clearance decreases over time, with a mean (CV%) maximal reduction from baseline to the 13^{th} weekly treatment dose of 40.8% (56%). The geometric mean (CV%) clearance is 0.472 L/day (64%) at the 13^{th} weekly treatment dose. Patients who discontinue teclistamab-cqyv after the 13^{th} weekly treatment dose are expected to have a 50% reduction from $C_{\rm max}$ in teclistamab-cqyv concentration at a median (5th to 95th percentile) time of 15 (7 to 33) days after $T_{\rm max}$ and a 97% reduction from $C_{\rm max}$ in teclistamab-cqyv concentration at a median time of 69 (32 to 163) days after $T_{\rm max}$.

Specific Populations

The volume of distribution and clearance of teclistamab-cqyv increase with increasing body weight (41 kg to 139 kg).

There were no clinically significant differences in the pharmacokinetics of teclistamab-cqyv based on age (24 to 84 years), sex, race (White, Black or African American), ethnicity (Hispanic/Latino, not Hispanic/Latino), mild or moderate renal impairment (estimated glomerular filtration rate [eGFR] by Modification of Diet in Renal Disease [MDRD] method: 30 to 89 mL/min), or mild hepatic impairment (total bilirubin less than or equal to upper limit of normal [ULN] with AST greater than ULN or total bilirubin greater than 1 to 1.5 times ULN with any AST). The effects of severe renal impairment (eGFR less than 30 mL/min) or moderate to severe hepatic impairment (total bilirubin greater than 1.5 times ULN with any AST) on the pharmacokinetics of teclistamab-cqyv are unknown.

Drug Interaction Studies

No clinical studies evaluating the drug interaction potential of teclistamab-cqyv have been conducted.

12.6 Immunogenicity

The observed incidence of anti-drug antibodies is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, including those of teclistamab-cqyv or of other teclistamab products.

During treatment in MajesTEC-1 (up to 27 months), 1/186 (0.5%) of patients treated with subcutaneous TECVAYLI at various dosages developed anti-teclistamab-cqyv antibodies. Because of the low occurrence of anti-drug antibodies, the effect of these antibodies on the pharmacokinetics, pharmacodynamics, safety, and/or effectiveness of teclistamab products is unknown.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No carcinogenicity or genotoxicity studies have been conducted with teclistamab-cqyv.

No studies have been conducted to evaluate the effects of teclistamab-cqyv on fertility.

14 CLINICAL STUDIES

The efficacy of TECVAYLI was evaluated in patients with relapsed or refractory multiple myeloma in a single-arm, open-label, multi-center study (MajesTEC-1, NCT03145181 [Phase 1] and NCT04557098 [Phase 2]). The study included patients who had previously received at least three prior therapies, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody. The study excluded patients who had stroke, seizure, allogeneic stem cell transplantation within the past 6 months, Eastern Cooperative Oncology Group (ECOG) performance score of 2 or higher, known active CNS involvement or clinical signs of meningeal involvement of multiple myeloma, or active or documented history of autoimmune disease, with the exception of vitiligo, Type 1 diabetes, and prior autoimmune thyroiditis.

Patients received step-up doses of 0.06 mg/kg and 0.3 mg/kg of TECVAYLI followed by TECVAYLI 1.5 mg/kg subcutaneously once weekly thereafter until disease progression or unacceptable toxicity [see Dosage and Administration (2.1)].

The efficacy population included 110 patients. The median age was 66 (range: 33 to 82) years with 16% of patients 75 years of age or older; 56% were male; 91% were White, 5% were Black or African American, 3% were Asian. The International Staging System (ISS) at study entry was Stage I in 50%, Stage II in 38%, and Stage III in 12% of patients. High-risk cytogenetics (presence of del(17p), t(4;14) and t(14;16)) were present in 25% of patients. Seventeen percent of patients had extramedullary plasmacytomas. Patients with prior BCMA-targeted therapy were not included in the efficacy population.

The median number of prior lines of therapy was 5 (range: 2 to 14); 78% of patients had received at least 4 prior lines of therapy. Eighty-one percent of patients received prior stem cell transplantation. All patients had received prior therapy with a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody, and 76% were triple-class refractory (refractory to a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody).

Efficacy was established based on overall response rate (ORR) as determined by the Independent Review Committee (IRC) assessment using International Myeloma Working Group (IMWG) 2016 criteria (see Table 13).

The median time to first response was 1.2 months (range: 0.2 to 5.5 months). With a median follow-up of 7.4 months among responders, the estimated duration of response (DOR) rate was 90.6% (95% CI: 80.3%, 95.7%) at 6 months and 66.5% (95% CI: 38.8%, 83.9%) at 9 months.

Table 13: Efficacy Results for MajesTEC-1

	N=110
Overall response rate (ORR: sCR+CR+VGPR+PR) n(%)	68 (61.8)
95% CI (%)	(52.1, 70.9)
Complete response (CR) or better ^a	31 (28.2)
Very good partial response (VGPR)	32 (29.1)
Partial response (PR)	5 (4.5)
Duration of Response (DOR) (months)	
DOR (Months): Median (95% CI)	NE (9.0, NE)

NE=not estimable

16 HOW SUPPLIED/STORAGE AND HANDLING

TECVAYLI® (teclistamab-cqyv) injection is a sterile, preservative-free, clear to slightly opalescent, colorless to light yellow solution supplied as follows:

- One 30 mg/3 mL (10 mg/mL) single-dose vial in a carton: NDC: 57894-449-01
- One 153 mg/1.7 mL (90 mg/mL) single-dose vial in a carton: NDC: 57894-450-01

Store refrigerated at 2 °C to 8 °C (36 °F to 46 °F) in the original carton to protect from light.

Do not freeze.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Cytokine Release Syndrome (CRS)

Discuss the signs and symptoms associated with CRS, including fever, hypoxia, chills, hypotension, sinus tachycardia, headache, and elevated liver enzymes. Advise patients to immediately contact their healthcare provider if they experience any signs or symptoms of CRS. Advise patients that they will be hospitalized for 48 hours after administration of all doses within the TECVAYLI step-up dosing schedule [see Dosage and Administration (2.4) and Warnings and Precautions (5.1)].

^a Complete response or better = Stringent complete response (sCR) + complete response (CR).

Neurologic Toxicity including ICANS

Discuss the signs and symptoms associated with neurologic toxicity, including ICANS, including headache, confusion, dysgraphia, motor dysfunction, neuropathy, or encephalopathy. Advise patients to immediately contact their healthcare provider if they experience any signs or symptoms of neurologic toxicity. Advise patients to refrain from driving or operating heavy or potentially dangerous machinery during and for 48 hours after completion of TECVAYLI step-up dosing schedule and in the event of new onset of any neurologic toxicity symptoms until neurologic toxicity resolves [see Dosage and Administration (2.4) and Warnings and Precautions (5.2)].

TECVAYLI and TALVEY REMS

TECVAYLI is available only through a restricted program called TECVAYLI and TALVEY REMS. Inform patients that they will be given a Patient Wallet Card that they should carry with them at all times and show to all of their healthcare providers. This card describes signs and symptoms of CRS and neurologic toxicity which, if experienced, should prompt the patient to immediately seek medical attention [see Warnings and Precautions (5.3)].

Hepatotoxicity

Advise patients that liver enzyme elevations may occur and that they should report symptoms that may indicate liver toxicity, including fatigue, anorexia, right upper abdominal discomfort, dark urine, or jaundice [see Warnings and Precautions (5.4)].

Infections

Discuss the signs and symptoms of infection [see Dosage and Administration (2.4) and Warnings and Precautions (5.5)].

Neutropenia

Discuss the signs and symptoms associated with neutropenia and febrile neutropenia [see Dosage and Administration (2.4) and Warnings and Precautions (5.6)]

Hypersensitivity and Other Administration Reactions

Advise patients to immediately seek medical attention for any signs and symptoms of systemic administration-related reactions. Advise patients that local injection-site reactions may occur and to report any severe reactions [see Warnings and Precautions (5.7)].

Embryo-Fetal Toxicity

Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to inform their healthcare provider if they are pregnant or become pregnant. Advise females of reproductive potential to use effective contraception during treatment with TECVAYLI and for 5 months after the last dose [see Warnings and Precautions (5.8) and Use in Specific Populations (8.1, 8.3)].

Lactation

Advise women not to breastfeed during treatment with TECVAYLI and for 5 months after the last dose [see Use in Specific Populations (8.2)].

Manufactured by: Janssen Biotech, Inc. Horsham, PA 19044, USA U.S. License Number 1864

For patent information: www.janssenpatents.com © Johnson & Johnson and its affiliates 2022

MEDICATION GUIDE TECVAYLI® [tek vay' lee]

(teclistamab-cgyv) injection, for subcutaneous use

What is the most important information I should know about TECVAYLI?

TECVAYLI may cause side effects that are serious, life-threatening or lead to death, including Cytokine Release Syndrome (CRS) and Neurologic problems.

Call your healthcare provider right away if you develop any of the signs or symptoms of CRS or neurologic problems listed below at any time during your treatment with TECVAYLI:

Cytokine Release Syndrome (CRS). Signs and symptoms of CRS may include:

• fever (100.4 °F or higher)

feeling anxious

difficulty breathing

· confusion or restlessness

chills

headache

· dizziness or lightheadedness

increased liver enzymes in your blood

· fast heartbeat

Neurologic problems. Symptoms of neurologic problems with TECVAYLI include:

headache

confusion

· changes in your handwriting

jerking movements

trouble speaking

· problems walking

· rigid muscles

muscle spasms

• muscle weakness in your body or face

feeling restless

tremor

hearing loss

• numbness and tingling (feeling like "pins and needles")

double vision

burning, throbbing, or stabbing pain

- Due to the risk of CRS and neurologic symptoms, you should be hospitalized for 48 hours after all doses of TECVAYLI that are part
 of the "step-up dosing schedule." The "step-up dosing schedule" is when you receive the first 2 doses of TECVAYLI, which are
 called "step-up" doses, and then you receive the first "treatment dose" of TECVAYLI. After "step-up" dose 1 of TECVAYLI, the
 dose of TECVAYLI is increased. After "step-up" dose 2, the dose is increased again when you receive the first "treatment dose" of
 TECVAYLI.
 - "Step-up dose 1" is given on day 1 of treatment. "Step-up dose 2" is usually given on day 4 of treatment. The first "treatment dose" is usually given on day 7 of treatment.
 - Your healthcare provider will decide when you will receive "step-up dose 2" and your first "treatment dose."
 - "Step-up" dose 2 may be given between 2 to 4 days after "step-up" dose 1, or up to 7 days after "step-up" dose 1 if you have certain side effects with TECVAYLI.
 - Your first "treatment dose" may be given between 2 to 4 days after "step-up" dose 2, or up to 7 days after "step-up" dose 2 if you have certain side effects with TECVAYLI.
 - Your healthcare provider will decide the number of days to wait between your doses of TECVAYLI as well as how many treatments you will receive.
 - If your dose of TECVAYLI is delayed for any reason, you may need to repeat the "step-up dosing schedule" to receive TECVAYLI.
 - Before each "step-up" dose and your first "treatment dose" of TECVAYLI you will receive medicines to help reduce your risk of CRS. Your healthcare provider will decide if you need to receive medicines to help reduce your risk of CRS with future doses.
- Your healthcare provider will monitor you for signs and symptoms of CRS and neurologic problems during treatment with TECVAYLI, as well as other side effects and treat you as needed.
- Do not drive or operate heavy or dangerous machinery during and for 48 hours after your TECVAYLI "step-up dosing schedule" is completed, or at any time during treatment with TECVAYLI if you develop new neurologic symptoms until the symptoms go away.

TECVAYLI is available only through the TECVAYLI and TALVEY Risk Evaluation and Mitigation Strategy (REMS) due to the risk of CRS and neurologic problems.

You will receive a Patient Wallet Card from your healthcare provider. Carry the Patient Wallet Card with you at all times and show it to all of your healthcare providers. The Patient Wallet Card lists signs and symptoms of CRS and neurologic problems.

Get medical help right away if you develop any of the signs and symptoms listed on the Patient Wallet Card. You may need to be treated in a hospital.

- If you have any questions about TECVAYLI, ask your healthcare provider.
- Your healthcare provider may temporarily stop or completely stop your treatment with TECVAYLI if you develop CRS, neurologic
 problems or any other side effects that are severe.

See "What are the possible side effects of TECVAYLI?" for more information about side effects.

What is TECVAYLI?

TECVAYLI is a prescription medicine to treat adults with multiple myeloma who:

- have already received at least 4 treatment regimens, including a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody to treat their multiple myeloma,
 and
- their cancer has come back or did not respond to prior treatment

It is not known if TECVAYLI is safe and effective in children.

Before you receive TECVAYLI, tell your healthcare provider about all of your medical conditions, including if you:

- · have an infection
- are pregnant or plan to become pregnant. TECVAYLI may harm your unborn baby.
 - Your healthcare provider should do a pregnancy test before you start treatment with TECVAYLI.
 - You should use effective birth control (contraception) during treatment and for 5 months after your last dose of TECVAYLI.
- Tell your healthcare provider right away if you become pregnant or think that you may be pregnant during treatment with TECVAYLI.
- are breastfeeding or plan to breastfeed. It is not known if TECVAYLI passes into your breast milk. Do not breastfeed during treatment and for 5 months after your last dose of TECVAYLI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will I receive TECVAYLI?

- TECVAYLI will be given to you by your healthcare provider as an injection under your skin (subcutaneous injection), usually in your stomach-area (abdomen), your thigh or another area of your body may be injected.
- See "What is the most important information I should know" at the beginning of this Medication Guide for information about how
 you will receive TECVAYLI.
- If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment. It is important for
 you to be monitored closely for side effects during treatment with TECVAYLI.

What are the possible side effects of TECVAYLI?

TECVAYLI may cause serious side effects, including:

- See "What is the most important information I should know about TECVAYLI?"
- Liver problems. TECVAYLI can cause liver problems that may lead to death. Increased bilirubin and liver enzymes in your blood are common with TECVAYLI and can also sometimes be severe These increases in liver enzymes can happen with or without you also having CRS. Your healthcare provider will monitor you for these problems before you start and during treatment with TECVAYLI.

 Tell your healthcare provider if you develop any symptoms of a liver problem including:
 - · tiredness
- pain in your right upper stomach-area (abdomen)
- loss of appetite
- · dark urine
- vellowing of your skin or white part of your eyes
- Infections. Upper respiratory tract infections and pneumonia are common with TECVAYLI. TECVAYLI can cause bacterial and viral infections that are severe, life-threatening, or that may lead to death.
 - Your healthcare provider will monitor you for signs and symptoms of infection before and during treatment with TECVAYLI.
 - Your healthcare provider may prescribe medicines for you to help prevent infections and treat you as needed if you develop an infection during treatment with TECVAYLI.
 - Tell your healthcare provider right away if you get a fever, chills or any signs or symptoms of an infection.
- Decreased white blood cell counts. Decreased white blood cell counts are common with TECVAYLI and can also be severe. Fever
 sometimes also happens with low white blood cell counts and may be a sign that you have an infection. Your healthcare provider
 will check your blood cell counts before you start and during treatment with TECVAYLI, and treat you as needed.
- Allergic reactions and injection site reactions. TECVAYLI can cause allergic reactions that can affect your whole body (systemic), and also cause injection site reactions.
 - Some people taking TECVAYLI can develop symptoms of an allergic reaction that can affect your whole body and may include fever or a swollen tongue. Get medical help right away if you develop symptoms of an allergic reaction during treatment with TECVAYLI.
 - Injection site reactions are common with TECVAYLI and can include: redness, heat, swelling, bruising, bacterial skin infection (cellulitis), discomfort, blood collection under the skin at the injection site (hematoma), and rash. Tell your healthcare provider if you develop any severe injection site reactions.

Your healthcare provider may temporarily or permanently stop TECVAYLI if you have any of the side effects listed above and they are severe.

The most common side effects of TECVAYLI include:

- fever
- pain in your joints and muscles, back and chest muscles, and in your arms and legs
- · tiredness and weakness

- upper respiratory tract infections and pneumonia.
 See "Infections" above.
- nausea
- headache
- diarrhea

The most common severe abnormal lab test results with TECVAYLI include: decreased white blood cells, red blood cells and platelets. These are not all the possible side effects of TECVAYLI.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General Information about TECVAYLI.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

You can ask your healthcare provider for information about TECVAYLI that is written for health professionals.

What are the ingredients of TECVAYLI?

Active ingredient: teclistamab-cqyv

Inactive ingredients: edetate disodium, glacial acetic acid, polysorbate 20, sodium acetate, sucrose, Water for Injection

Manufactured by: Janssen Biotech, Inc., Horsham, PA 19044, USA

U.S. License Number 1864

For patent information: www.janssenpatents.com

© Johnson & Johnson and its affiliates 2022

For more information about TECVAYLI go to www.TECVAYLI.com or call 1-800-526-7736.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: Aug 2025

cp-326438v12